

Marine Mammals: Multiple Small Stocks with Insufficient Data-Bay, Sound, and Estuary Stocks

NOAAFISHERIES


Southeast Fisheries Science Center

SEFSC Protected Species Program Review

25-27 August 2015 Miami, Florida

- Stocks first delimited in 1994-1995 with the first Stock Assessment Reports
 - Originally 33 Bay, Sound and Estuary stocks (now 32)
 - Small areas, low abundance
 - Experience multiple lethal and sub-lethal impacts

Challenge for research and management






- Progression in Gulf of Mexico- Abundance
 - 24 stocks (77%) have abundance estimates > 8 years old
 - Standard aerial line-transect surveys generally not appropriate

Sought to standardize and promote consistent use of photo-ID capture mark-recapture methodologies





<u>Critical Methodological Considerations</u>

- Survey design
 - Spatial extent
 - Transients and seasonal timing
 - Recommend "Robust-design"
- Photo quality scoring and photo-id analysis procedures



NOAA Technical Memorandum NMFS-SEFSC-621

PHOTO-IDENTIFICATION CAPTURE-MARK-RECAPTURE TECHNIQUES FOR ESTIMATING ABUNDANCE OF BAY, SOUND AND ESTUARY POPULATIONS OF BOTTLENOSE DOLPHINS ALONG THE U.S. EAST COAST AND GULF OF MEXICO: A WORKSHOP REPORT

BY


PATRICIA E. ROSEL, KEITH D. MULLIN, LANCE GARRISON, LORI SCHWACKE, JEFF ADAMS, BRIAN BALMER, PAUL CONN, MICHAEL J. CONROY, TOMO BUUCHI, ANNIE GORGONE, ALETA HOHN, MARILYN MAZZOIL, CARL SCHWARZ, CARRIE SINCLAIR, TODD SPEAKMAN, KIM URIAN, NICOLE VOLLMER, PAUL WADE, RANDALL WELLS, ERIC ZOLMAN



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Flaharies Service
Southeast Fisheries Science Center
646 Cajundeme Boulevard
Lafavette, LA 7056 USA

September 201

 Progression in Gulf of Mexico - Abundance estimates less than 8 years old (n=9)





 Progression in Gulf of Mexico – Stock Structure · Complicated by proximity of bays, potertial of ba transients Possibility of population structure within some of the Bays ve 3 Texas BSE
Stocks
Sample a Bayrang adjacent coastal waters • Surve 3 Texas BSE 30 m season when transients are least likely **Christi Bay** Gulf of Mexico 25 50 Km




- Progression in Gulf of Mexico Stock Structure
 - First confirmation of demographic independence of BSE stocks achieved through genetic analysis (Sellas *et al.* 2005)






Bay, Sound and Estuary Stocks - Atlantic

- No BSE stocks delimited in the Atlantic with the first Stock Assessment Reports in 1994-1995
- 1996-present: Variety of studies indicate resident stocks
 - Photo-ID
 - Habitat inference for boundaries
 - Genetics
 - Stable isotopes
 - Telemetry






Atlantic Estuarine Year-Round Resident Stocks











Bay, Sound and Estuary Stocks - Mortality

- Most stocks do not have mortality estimates
- Fishery mortality does occur
 - Inshore shrimp trawl fishery is not observed
 - Crab/trap pot fisheries, commercial and recreational, unobserved
 - Mid-Atlantic gillnet fishery part of TRT for Atlantic stocks, observed
 - Documented takes are known to occur but are very rare- e.g.,
 1 observed take between 2007-2011. But stranding data suggest more interactions
 - Observer coverage is 1-3%
 - Very difficult to obtain accurate fishery mortality estimates
- Variety of other anthropogenic impacts (contaminants, habitat loss)




Challenges

- Estuarine bottlenose dolphins have complicated stock structure
 - Exhibit high site fidelity
 - Genetic data to date indicate limited gene flow
 - Represent demographically independent populations
 - Multiple lethal and sub-lethal stressors
- Requires dedicated crews for photo-ID to keep up with abundance estimate requirements
- Requires biopsy effort and stock structure research program to ensure stocks are correctly delineated
- Capacity allows for one CMR or biopsy survey per year
- Need for prioritization



Bay, Sound and Estuary Stocks - GOMx: Prioritization of Stocks





Bay, Sound and Estuary Stocks - GOMx: Prioritization of Research



NOAA Technical Memorandum NMFS-SEFSC-665

A Method for Prioritizing Research on Common Bottlenose Dolphin Stocks
Through Evaluating Threats and Data Availability:
Development and Application to Bay, Sound and Estuary Stocks in Texas

BY NICOLE M. PHILLIPS AND PATRICIA E. ROSEL



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southeast Fisheries Science Center
646 Cajundome Boulevard
Lafayette, LA 70506 USA

December 2014

- Developed a framework to prioritize BSE stocks for stock assessment research
- Accounts for potential impact of suite of stressors and for availability of stock assessment data
- Sets priority levels based upon threshold values

Stressors

Habitat loss

Adverse weather Freshwater inflows

Freshwater inflows

Shipping

Tourism & boat traffic

of unknown etiology

Unusual mortality events

Heavy metal pollution

Chemical pollution Hypoxia

Climate change Marine debris

Commercial fisheries Noise

Algal blooms


Aquaculture

Disease Oil & gas pollution

Dredging & construction Recreational fisheries



Bay, Sound and Estuary Stocks - GOMx: Prioritization for Research



Cumulative Threat Score

Based on the potential impacts of the 19 anthropogenic and environmental stressors

Data Availability Score

Reflects information about stock structure, abundance, and mortality

High priority stocks have high potential threats and low data availability

Texas stocks completed Louisiana stocks in progress



Strengths

- Provide standardized tools/methods to external partners for abundance estimates
- Plan for prioritizing stocks
- Proficiency with biopsy sampling and capture markrecapture photo-ID field work and photo analysis
- Strong stock structure investigation program



Future Directions

- Complete research prioritization for Gulf of Mexico BSE stocks and adapt prioritization process to Atlantic BSE stocks, oceanic stocks
- Conduct GOMx-wide genetic analysis of BSE stock structure with samples available to date
- Ecosystem-based approach: Investigate alternative models to incorporate more than just commercial fishery interactions as the metric for status of a stock; risk assessment that incorporate status and trends, identify major risks, quantify the cumulative impacts of human activities and environmental variation
- Improve demographic population models that incorporate reproductive parameters (which we don't yet have)



Discussion Topics

- Is the work we are doing reflective of scientific best practices?
- Do you see an opportunity for SEFSC to shift resources from an existing activity to deal with an unmet need?
- Are stocks being prioritized properly?
- Discuss the major limitations/weaknesses of protected species abundance, distribution and genetic studies and how could they be resolved?

